

NEW RESULTS ON HYPOHAMILTONIAN AND ALMOST HYPOHAMILTONIAN GRAPHS

JAN GOEDGEBEUR* AND CAROL T. ZAMFIRESCU*

Abstract. Consider a non-hamiltonian graph G . G is *hypohamiltonian* if for every vertex v in G , the graph $G - v$ is hamiltonian. G is *almost hypohamiltonian* if there exists a vertex w in G such that $G - w$ is non-hamiltonian, and $G - v$ is hamiltonian for every vertex $v \neq w$. McKay asked in [*J. Graph Theory*, doi: 10.1002/jgt.22043] whether infinitely many planar cubic hypohamiltonian graphs of girth 5 exist. We settle this question affirmatively.

The second author asked in [*J. Graph Theory* **79** (2015) 63–81] to find all orders for which almost hypohamiltonian graphs exist—we solve this problem as well. To this end, we present a specialised algorithm which generates complete sets of almost hypohamiltonian graphs for various orders. Furthermore, we show that the smallest cubic almost hypohamiltonian graphs have order 26. We also provide a lower bound for the order of the smallest planar almost hypohamiltonian graph and improve the upper bound for the order of the smallest planar almost hypohamiltonian graph containing a cubic vertex. Finally, we determine the smallest planar almost hypohamiltonian graphs of girth 5, both in the general and cubic case.

Key words. Hamiltonian, hypohamiltonian, almost hypohamiltonian, planar, cubic, exhaustive generation

AMS subject classifications. 05C10, 05C35, 05C38, 05C45, 05C85

1. Introduction. Every graph in this paper will be undirected, finite, connected, and neither contain loops nor multiple edges, unless stated otherwise. A graph is *hamiltonian* if it contains a cycle visiting every vertex of the graph—such a cycle is called *hamiltonian*. A graph G is *hypohamiltonian* if G itself is non-hamiltonian, but for every vertex v in G , the graph $G - v$ is hamiltonian. The study of hypohamiltonicity goes back to a paper of Sousselier [25] from the early sixties. The survey of Holton and Sheehan [20] provides a good overview. New material published after the appearance of this survey includes an article of Jooyandeh, McKay, Östergård, Pettersson, and the second author [22], work of McKay [23], and the paper [17] by the two authors of this work.

We now introduce our terminology—see [13] for undefined notions. We call a vertex *cubic* if it has degree 3, and a graph *cubic* if all of its vertices are cubic. Consider a graph G . Two edges of G are *independent* if they have no common vertices. We use $\deg(v)$ to denote the degree of a vertex v and put $\Delta(G) = \max_{v \in V(G)} \deg(v)$. The *girth* of a graph is the length of its shortest cycle. A cycle of length k will be called a *k-cycle*. For $S \subset V(G)$, $G[S]$ shall denote the graph induced by S . A not necessarily connected subgraph $G' = (V', E') \subset G = (V, E)$ is *spanning* if $V' = V$. For a set X , we denote by $|X|$ its cardinality. Following Aldred, McKay, and Wormald [2], we say that a graph G is *hypocyclic* if for every $v \in V(G)$, the graph $G - v$ is hamiltonian. Hamiltonian hypocyclic graphs are called “1-hamiltonian”, see e.g. [10], so the family of all hypocyclic graphs is the disjoint union of the families of all 1-hamiltonian and hypohamiltonian graphs. A graph G is *almost hypocyclic* if there exists a $w \in V(G)$ such that $G - w$ is non-hamiltonian, but for every vertex $v \neq w$ in G , the graph $G - v$ is hamiltonian. We call w the *exceptional vertex* of G . If G is additionally non-

*Department of Applied Mathematics, Computer Science & Statistics, Ghent University, Krijgslaan 281-S9, 9000 Ghent, Belgium (jan.goedgebeur@ugent.be, czamfirescu@gmail.com). The first author is supported by a Postdoctoral Fellowship of the Research Foundation Flanders (FWO). The second author is supported by a PhD fellowship at Ghent University on the Special Research Fund (BOF) scholarship 01DI1015.

hamiltonian, then G is called *almost hypohamiltonian* (with exceptional vertex w).

This article is structured as follows. In Section 2 we solve a problem of McKay: he asked [23] whether infinitely many planar cubic hypohamiltonian graphs of girth 5 exist. We settle this question affirmatively.

In Section 3 we deal with almost hypohamiltonian graphs, extending a series of results (and settling certain open questions) of the second author [32, 33]. In particular, in Section 3.1 we present a specialised algorithm for generating all almost hypohamiltonian graphs. We also use our implementation of this algorithm to generate complete sets of almost hypohamiltonian graphs for various orders and in Section 3.2 we show that there exists an almost hypohamiltonian graph of order n iff $n \geq 17$, solving [32, Problem 2].

In Section 3.3 we show that the smallest cubic almost hypohamiltonian graphs have order 26, which settles the non-planar variant of [32, Problem 1]—the planar version had been solved by McKay (private communication). In Section 3.4 we give a lower bound for the order of the smallest planar almost hypohamiltonian graphs, and improve the upper bound for the order of the smallest planar almost hypohamiltonian graph whose exceptional vertex is cubic (from 47, see [32], to 36), which are particularly important due to results from [32]. This answers [33, Problem 6(a)] and [33, Problem 7]. We also prove that such a graph of order n exists for every $n \geq 73$, partially answering [33, Problem 6(b)]. Furthermore, we show that the smallest planar almost hypohamiltonian graph of girth 5 has order 44. (In the hypohamiltonian case, this number is 45, see [22].) Finally, in Section 3.5 we determine that the smallest planar cubic almost hypohamiltonian graph has order at least 54, and that the smallest such graph which additionally has girth 5, is of order 68.

2. McKay’s Problem on planar cubic hypohamiltonian graphs. In [23], McKay constructed three planar cubic hypohamiltonian graphs of girth 5 and raised the natural question whether infinitely many such graphs exist. In this section we prove that this question has a positive answer. Let G be a graph containing vertices v, w, x, y . Following Chvátal [11], the pair of vertices (v, w) is *good* in G if there exists a hamiltonian path in G with end-vertices v and w . Two pairs of vertices $((v, w), (x, y))$ are *good* in G if there exist two disjoint paths which together span G , and which have end-vertices v and w , and x and y , respectively.

Let H and H' be cubic graphs on at least 6 vertices. Consider $J = H - \{ab, cd\}$, where ab and cd are independent edges in H , $J' = H' - \{x, y\}$, where x and y are adjacent vertices in H' , and let a', b' and c', d' be the other neighbours of x and y in H' , respectively. Then the *dot product* $H \cdot H'$ is defined as the graph

$$(V(H) \cup V(J'), E(J) \cup E(J') \cup \{aa', bb', cc', dd'\}).$$

(Note that under above conditions, the dot product may be disconnected. The dot product was defined by Isaacs in [21]. In fact, following Skupień [24], it seems that this operation was invented earlier by Adel’son-Vel’skiĭ and Titov [1].)

THEOREM 1. *Let H and H' be cubic graphs, H non-hamiltonian and H' hypohamiltonian. Consider independent edges $ab, cd \in E(H)$. If each of*

$$(a, c), (a, d), (b, c), (b, d), \text{ and } ((a, b), (c, d))$$

is good in $H - \{ab, cd\}$, and for each vertex $t \in V(H)$, at least one of (a, b) or (c, d) is good in $H - \{t, ab, cd\}$, then $H \cdot H'$ is a cubic hypohamiltonian graph as well. If H and H' are planar, and ab and cd lie on the same facial cycle, then the dot product

can be applied such that $H \cdot H'$ is planar, as well. If g (g') is the girth of H (H'), then the girth of $H \cdot H'$ is at least $\min\{g, g'\}$.

Proof. Put $G = H \cdot H'$ and denote by x, y the (adjacent) vertices which were deleted from H' to form the dot product. G is obviously cubic. Put $N(x) = \{a', b', y\}$ and $N(y) = \{c', d', x\}$ such that the unique neighbour of a' (b' , c' , d') in G is a (b , c , d). We treat $H - \{ab, cd\} = J$ and $H' - \{x, y\} = J'$ as subgraphs of G . W.l.o.g. we may assume that the aforementioned labeling of vertices satisfies that if H and H' are planar, then G is also planar.

We first show that G is non-hamiltonian. Assume there exists a hamiltonian cycle $\mathfrak{h}$ in G . If $\mathfrak{h} \cap J$ consists of two components P and P' , then either $\mathfrak{h} \cap J + ab + cd$ is a hamiltonian cycle in H , a contradiction since H is non-hamiltonian, or P is a path with end-vertices a and b , and P' is a path with end-vertices c and d . But then $\mathfrak{h} \cap J'$ consists of two disjoint paths which together with $a'xb'$ and $c'yd'$ form a hamiltonian cycle in H' , a contradiction since H' is hypohamiltonian. So $\mathfrak{h} \cap J$ consists of one component. If $\mathfrak{h} \cap J$ has end-vertices a, b or c, d , then we immediately obtain a contradiction (by adding the edge ab or cd) to the non-hamiltonicity of H , so we may assume that $\mathfrak{h} \cap J'$ is a hamiltonian path P with end-vertices $p \in \{a', b'\}$ and $q \in \{c', d'\}$. Then $P \cup pxyq$ is a hamiltonian cycle in H' , once again a contradiction.

It remains to show that $G - s$ is hypocyclic for every $s \in V(G)$. Let $s \in V(J')$. Since H' is hypohamiltonian, there exists a hamiltonian cycle $\mathfrak{h}$ in $H' - s$. Put $Q = a'xb'$ and $Q' = c'yd'$. Note that if exactly one of Q and Q' lies in $\mathfrak{h}$, then necessarily $s \in \{x, y\}$, which is impossible, as neither x nor y lie in J' .

Case 1: Neither Q nor Q' lie in $\mathfrak{h}$. In this situation, $xy \in E(\mathfrak{h})$, and say a' and d' are the end-vertices of the path $P' = \mathfrak{h} - \{x, y\}$, which is a hamiltonian path in J' . (a, d) is good in J , i.e. there exists a hamiltonian path P in J with end-vertices a, d , so we obtain the hamiltonian cycle $P' \cup P + aa' + dd'$ in $G - s$ as desired. (If P' has end-vertices other than a', d' , the argument is very similar. Likewise if $s \in \{a', b', c', d'\}$.)

Case 2: Both Q and Q' lie in $\mathfrak{h}$. $((a, b), (c, d))$ is good in H , so there exist disjoint paths P_1, P_2 in H which together span H , with end-vertices a, b and c, d , respectively. Denote by P'_1, P'_2 the components of $\mathfrak{h} - \{x, y\}$. Then $P_1 \cup P_2 \cup P'_1 \cup P'_2 + aa' + bb' + cc' + dd'$ is a hamiltonian cycle in $G - s$.

Now consider $s \in V(H)$. We know that at least one of (a, b) and (c, d) is good in $H - \{s, ab, cd\}$, say (a, b) . So there exists a hamiltonian path P in $H - \{s, ab, cd\}$ with end-vertices a and b . Since H' is hypohamiltonian, $H' - y$ is hamiltonian, which implies that there exists a hamiltonian path P' in J' with end-vertices a' and b' . Now $P \cup P' + aa' + bb'$ is a hamiltonian cycle in $G - s$ (here it is crucial that $cd \notin E(P)$, as $cd \in E(H)$, but $cd \notin E(G)$). If (c, d) is good in $H - \{s, ab, cd\}$ and not (a, b) , we use the fact that $H' - x$ is hamiltonian.)

Finally, let g be the girth of H and g' be the girth of H' . Assume the girth of $H \cdot H'$ is $k < \min\{g, g'\}$, and let C be a k -cycle in $H \cdot H'$. Clearly, C contains (at least) two edges $e_1, e_2 \in \{aa', bb', cc', dd'\}$ with $e_1 = aa'$ w.l.o.g. If $e_2 = bb'$, then $(C \cap J') \cup a'xb'$ is a cycle of length at most $k - 2$ in H' . If $e_2 = cc'$, then $(C \cap J') \cup a'xy c'$ is a cycle of length at most k in H' . If $k < g'$, in both cases we have a contradiction, so $k \geq g'$. But this is impossible, as $k < \min\{g, g'\}$. We have shown that the girth of $H \cdot H'$ is at least $\min\{g, g'\}$. $\square$

The above result is inspired by a theorem of Fiorini [14], who showed that gluing two hypohamiltonian snarks using the dot product yields a new hypohamiltonian

snark. Note that in Fiorini's statement the hypotheses he formulates are too weak to prove the result he advertises; Cavicchioli et al. [9] have identified this omission, but we believe their version is also not entirely correct. A detailed account will be given in the note [15].

In the following, we will call the pair of edges ab, cd from the statement of Theorem 1 *suitable*. Let H be the graph from Figure 1. It is one of the three planar cubic hypohamiltonian graphs of girth 5 on 76 vertices found by McKay [23]. Using a computer, we found several suitable pairs of edges ab, cd in this graph. Since we will use double-digit numbers to label the vertices, we shall denote the edge vw by (v, w) for better readability.




Fig. 1: The planar cubic hypohamiltonian graph H . It has girth 5 and order 76. H is one of the three smallest planar cubic hypohamiltonian graphs of girth 5, and was found by McKay [23]. The suitable edge pair $(20, 28), (52, 62)$ is marked in bold.

LEMMA 2. Let H be the graph from Figure 1. The edge pair $(20, 28), (52, 62)$ is suitable in H .

Proof. It is clear from Figure 1 that the edges $(20, 28)$ and $(52, 62)$ are on the same facial cycle. (This guarantees that the dot product is planar.)

$(20, 52), (20, 62), (28, 52)$ and $(28, 64)$ are good in $H - \{(20, 28), (52, 62)\}$ due to

the following hamiltonian paths, respectively:

- 52, 44, 36, 26, 16, 8, 3, 7, 14, 22, 23, 15, 24, 25, 34, 41, 42, 35, 43, 51, 61, 71, 72, 62, 53, 45, 37, 29, 30, 38, 46, 55, 54, 64, 63, 73, 76, 70, 60, 69, 75, 74, 66, 65, 56, 47, 57, 67, 68, 59, 50, 58, 49, 48, 40, 33, 32, 39, 31, 21, 13, 6, 2, 1, 4, 9, 17, 27, 28, 19, 18, 10, 5, 11, 12, 20
- 62, 53, 45, 37, 29, 30, 38, 46, 55, 54, 64, 63, 73, 72, 71, 61, 52, 44, 36, 26, 35, 43, 51, 60, 70, 76, 75, 69, 59, 68, 74, 66, 65, 56, 47, 57, 67, 58, 50, 42, 41, 49, 48, 39, 32, 23, 15, 24, 33, 40, 34, 25, 16, 8, 3, 7, 14, 22, 31, 21, 13, 6, 2, 1, 4, 9, 17, 27, 28, 19, 18, 10, 5, 11, 12, 20
- 52, 44, 36, 26, 16, 25, 24, 15, 7, 14, 13, 21, 30, 38, 46, 55, 54, 64, 63, 73, 76, 70, 60, 69, 75, 74, 66, 65, 56, 47, 57, 67, 68, 59, 50, 58, 49, 48, 39, 31, 22, 23, 32, 33, 40, 34, 41, 42, 35, 43, 51, 61, 71, 72, 62, 53, 45, 37, 29, 20, 12, 6, 2, 1, 3, 8, 9, 4, 10, 5, 11, 19, 18, 17, 27, 28
- 62, 53, 45, 37, 29, 20, 12, 6, 13, 14, 7, 15, 23, 22, 31, 21, 30, 38, 46, 55, 54, 64, 63, 73, 72, 71, 70, 76, 75, 69, 60, 51, 61, 52, 44, 43, 35, 42, 50, 59, 68, 74, 66, 65, 56, 47, 57, 67, 58, 49, 41, 34, 40, 48, 39, 32, 33, 24, 25, 16, 26, 36, 27, 17, 18, 10, 4, 9, 8, 3, 1, 2, 5, 11, 19, 28

$((20, 28), (52, 62))$ is good in $H - \{(20, 28), (52, 62)\}$ due to the following two disjoint paths with end-vertices 20 and 28, and 52 and 62, respectively, which together span H .

- 28, 19, 11, 12, 20
- 62, 53, 45, 37, 29, 30, 21, 13, 6, 2, 5, 10, 18, 17, 27, 36, 26, 35, 42, 41, 34, 25, 16, 8, 9, 4, 1, 3, 7, 14, 22, 31, 39, 32, 23, 15, 24, 33, 40, 48, 49, 58, 50, 59, 69, 60, 70, 76, 75, 74, 68, 67, 57, 66, 65, 56, 47, 38, 46, 55, 54, 64, 63, 73, 72, 71, 61, 51, 43, 44, 52

The fact that at least one of $(20, 28)$ or $(52, 62)$ is good in $H - \{t, (20, 28), (52, 62)\}$ for every $t \in V(H)$ was shown by computer. We do not include the 76 hamiltonian paths here due to space constraints—76 paths are necessary since H has trivial automorphism group [23]—, but in each case we verified that the path found by the computer is indeed a valid hamiltonian path in the graph. $\square$

As in Lemma 2, H will denote in the following the 76-vertex graph from Figure 1. Consider a copy H' of H . Using the pair of suitable edges lying on the same facial cycle in H we have found in Lemma 2, and any pair of adjacent vertices in H' , we form the graph $H \cdot H'$. By Theorem 1, $H \cdot H'$ is a planar cubic hypohamiltonian graph of girth 5. (Theorem 1 states that the girth of $H \cdot H'$ is at least 5, and a planar 3-connected graph cannot have girth greater than 5.) Iterating this process (i.e., for a copy H'' of H , forming the dot product $H'' \cdot (H \cdot H')$, etc.) we obtain the following answer to McKay’s question [23]. The second part of the statement was obtained by using the 78-vertex planar cubic hypohamiltonian graph of girth 5 constructed in [17] as H' in Theorem 1.

THEOREM 3. *There are infinitely many planar cubic hypohamiltonian graphs of girth 5. More specifically, there exist such graphs of order n for every $n = 74k + a$, where $k \geq 2$ and $a \in \{2, 4\}$.*

We remark that Theorem 1 may in fact be applied to show that there are smaller graphs which could be used in the above construction, since in its statement we only ask of H to be non-hamiltonian, not hypohamiltonian. However, our focus here was to solve McKay’s problem—future work may include a more careful analysis of the graphs which may take the place of H .

We shall return to planar 3-connected cubic graphs with interesting hypohamiltonian properties at the end of this paper.

3. Almost hypohamiltonian graphs.

3.1. The generation algorithm. In [17] we presented an algorithm for the exhaustive generation of hypohamiltonian graphs. It is based on work of Aldred, McKay, and Wormald [2]. We now describe how we modified this algorithm to generate almost hypohamiltonian graphs exhaustively. We point out that in the following arguments there is significant overlap with [17]—still, we choose to present these results for two reasons: firstly, small changes with respect to [17] are nonetheless necessary and important, and secondly, we would like to make this paper to a large extent self-contained.

Let G be a possibly disconnected graph, and $p(G)$ the minimum number of disjoint paths needed to cover all vertices of G . Denote by $V_1(G)$ the vertices of degree 1 in G , and by $I(G)$ the set of all isolated vertices and all isolated edges of G . Put

$$k(G) = \begin{cases} 0 & \text{if } G \text{ is empty,} \\ \max \left\{ 1, \left\lceil \frac{|V_1|}{2} \right\rceil \right\} & \text{if } I(G) = \emptyset \text{ but } G \text{ is not empty,} \\ |I(G)| + k(G - I(G)) & \text{else.} \end{cases}$$

We now adapt two lemmas of Aldred, McKay, and Wormald [2] from the hypocyclic to the almost hypocyclic case.

LEMMA 4. *Given an almost hypocyclic graph G with exceptional vertex w , for any partition (W, X) of the vertices of G with $|W| > 1$ and $|X| > 1$, we have that*

$$p(G[W]) < |X| \quad \text{and} \quad k(G[W]) < |X|.$$

Consider a graph G containing a partition (W, X) of its vertices with $|W| > 1$ and $|X| > 1$. If $p(G[W]) \geq |X|$, then we call (W, X) a *type A obstruction*, and if $k(G[W]) \geq |X|$, then we speak of a *type B obstruction*. For efficiency reasons we only consider type A obstructions where $G[W]$ is a union of disjoint paths in the algorithm.

LEMMA 5. *Let G be an almost hypocyclic graph with exceptional vertex w , and consider a partition (W, X) of the vertices of G with $|W| > 1$ and $|X| > 1$ such that W is an independent set. Furthermore, for some vertex $v \in X$, define n_1 and n_2 to be the number of vertices of $X - v$ joined to one or more than one vertex of W , respectively. Then we have $2n_2 + n_1 \geq 2|W|$ for every $v \in X$.*

If all assumptions of Lemma 5 are met and $2n_2 + n_1 < 2|W|$ for some $v \in X$, we call (W, X, v) a *type C obstruction*. The following proof is a straightforward adaptation of the proof of Aldred, McKay, and Wormald [2].

Proof of Lemmas 4 and 5. Consider $v \in X$ with $v \neq w$ (this choice is possible, since $|X| > 1$), and let $\mathfrak{h}$ be a hamiltonian cycle in $G - v$. The number of components of $\mathfrak{h}$ restricted to W must be at least $p(G[W])$, so the number of components in $\mathfrak{h}$ restricted to $X - v$ is at least $p(G[W])$ (and thus we have $|X| - 1 \geq p(G[W])$). This handles type A and, since clearly $k(G) \leq p(G)$, type B obstructions. For type C, consider the same cycle $\mathfrak{h}$. The number of edges of $\mathfrak{h}$ between W and X must be $2|W|$, but X can supply at most $2n_2 + n_1$. $\square$

LEMMA 6. *Let G be an almost hypohamiltonian graph with exceptional vertex w . Then G is 3-connected. In particular, G has minimum degree 3.*

Proof. Since for every vertex $v \neq w$, the graph $G - v$ is hamiltonian, G must be 2-connected. Now let $\{x, y\}$ be a 2-cut in G . Both $G - x$ and $G - y$ are non-hamiltonian, which is impossible. So G does not contain 2-cuts, thus G is 3-connected. $\square$

Note that if we allow more than one exceptional vertex, 3-connectedness is not guaranteed anymore. Or put in rigorous terms: consider a 2-connected graph G of circumference $|V(G)| - 1$ and let $W \subset V(G)$ be the (possibly empty) inclusion-maximal set of vertices such that for all $w \in W$ the graph $G - w$ is non-hamiltonian. (And thus, for all $v \in V(G) \setminus W$, the graph $G - v$ is hamiltonian.) We say that G is $|W|$ -hypohamiltonian. Although 0-hypohamiltonian (i.e. hypohamiltonian) and 1-hypohamiltonian (i.e. almost hypohamiltonian) graphs are 3-connected, it is easy to see that 2-hypohamiltonian graphs of connectivity 2 exist: consider a 4-cycle $v_1 v_2 v_3 v_4$, add an extra vertex v_5 , and the edges $v_2 v_5$ and $v_4 v_5$. This emphasises the intrinsic qualitative difference between $\{0, 1\}$ -hypohamiltonian graphs and k -hypohamiltonian graphs with $k \geq 2$.

LEMMA 7. *Let M be a 3-edge-cut in an almost hypohamiltonian graph G with exceptional vertex w . Then $G - M$ contains exactly two components A_1 and A_2 with $A = K_1$ and $A_2 \neq K_1$. In particular, almost hypohamiltonian graphs are cyclically 4-edge-connected.*

Proof. Let $G - M$ contain components $A_1, \dots, A_k$, $k \geq 2$, and assume that $A_1 \neq K_1$ and $A_2 \neq K_1$. We put $M = \{a_1 b_1, a_2 b_2, a_3 b_3\}$, where $a_i \in V(A_1)$, $b_i \in V(A_2)$ for all i . Since G is 3-connected, the elements of the set $\{a_1, a_2, a_3, b_1, b_2, b_3\}$ are pairwise distinct. Since G is almost hypohamiltonian, there exists an i such that $G - b_i$ and $G - a_i$ both are hamiltonian. Hence, there is a hamiltonian path in A_1 with end-vertices a_j and a_k , $j \neq i \neq k$, and a hamiltonian path in A_2 with end-vertices b_j and b_k , $j \neq i \neq k$. These paths together with $a_j b_j$ and $a_k b_k$ yield a hamiltonian cycle in G , a contradiction. Note that if at least one of the components of $G - M$ is isomorphic to K_1 , then $k = 2$. This completes the proof. $\square$

Collier and Schmeichel [12, p. 196] observed that the vertices of a triangle in a hypohamiltonian graph have degree at least 4. We now show that this holds for almost hypohamiltonian graphs, as well.

LEMMA 8. *Let G be an almost hypohamiltonian graph containing a triangle T . Then every vertex of T has degree at least 4.*

Proof. Put $V(T) = \{v_1, v_2, v_3\}$, where v_3 shall be cubic. Since G is almost hypohamiltonian, at least one of $G - v_1$ and $G - v_2$ must be hamiltonian, say $G - v_1$. Let $\mathfrak{h}$ be a hamiltonian cycle in $G - v_1$. As $v_3 v_2 \in E(\mathfrak{h})$, replace in $\mathfrak{h}$ the edge $v_3 v_2$ with the path $v_3 v_1 v_2$. We obtain a hamiltonian cycle in G , a contradiction. $\square$

We make use of these lemmas in our algorithm to generate all almost hypohamiltonian graphs—first we give an intuitive description, which is followed by the rigorous argument. Intuitively, by a *good Y-edge* we mean an edge which works towards the removal of a type Y obstruction, where $Y \in \{A, B, C\}$. Let us now formally define these good Y -edges.

Given a hypohamiltonian graph G' , and let G be a spanning subgraph of G' which contains a type A obstruction (W, X) (where $G[W]$ is a union of disjoint paths), since G' is hypohamiltonian it cannot contain a type A obstruction. Thus, there is an edge $e \in E(G') \setminus E(G)$ the endpoints of which lie in different components of $G[W]$. Furthermore, at least one of the endpoints of e has degree at most one in $G[W]$. We call such an edge a *good A-edge* for (W, X) . In the same vein, a *good B-edge* for a

type B obstruction (W, X) in G is a non-edge of G that joins two vertices of W , where at least one of those vertices has degree at most one in $G[W]$. Finally, a *good C-edge* for a type C obstruction (W, X, v) in G is a non-edge e of G for which one of the two following conditions holds:

- (i) Both endpoints of e are in W .
- (ii) One endpoint of e is in W and the other endpoint is in $X - v$ and has at most one neighbour in W .

It is straightforward that this is the only way to destroy a type B/C obstruction.

The pseudocode of our enumeration algorithm is given in Algorithm 1 and Algorithm 2. It is analogous to the algorithm we presented in [17] to generate all hypohamiltonian graphs. Nonetheless, we choose to present it here as well for the sake of convenience.

In order to generate all almost hypohamiltonian graphs with n vertices we start from a graph G which consists of an $(n - 1)$ -cycle and an isolated vertex h (disjoint from the cycle), so $G - h$ is hamiltonian. We then connect h to D vertices of the $(n - 1)$ -cycle in all possible ways and then perform Algorithm 2 on these graphs. Algorithm 2 will continue to recursively add edges in all possible ways such that at most one vertex w , the exceptional vertex, has degree larger than D .

It is essential for the efficiency of the algorithm that as few as possible edges are added by Algorithm 2, while still guaranteeing that all almost hypohamiltonian graphs are found by the algorithm.

We omit the proof of the following theorem as it is analogous to the proof of [17, Theorem 2.8].

THEOREM 9. *If Algorithm 1 terminates, the list of graphs $\mathcal{H}$ outputted by the algorithm is the list of all almost hypohamiltonian graphs with n vertices.*

Algorithm 1 Generate all almost hypohamiltonian graphs with n vertices

```

1: let  $\mathcal{H}$  be an empty list
2: let  $G := C_{n-1} + h$ 
3: for all  $3 \leq D \leq n - 1$  do
4:   // Generate all almost hypohamiltonian graphs where at least one vertex has
   // degree  $D$  and at most one vertex has degree larger than  $D$ 
5:   for every way of connecting  $h$  of  $G$  with  $D$  vertices of the  $C_{n-1}$  do
6:     Call the resulting graph  $G'$ 
7:     Construct( $G', D$ ) // i.e. perform Algorithm 2
8:   end for
9: end for
10: Output  $\mathcal{H}$ 

```

Note that since our algorithm only adds edges and never removes any vertices or edges, all graphs obtained by the algorithm from a graph with a g -cycle will have a cycle of length at most g . So in case we only want to generate almost hypohamiltonian graphs with a given lower bound k on the girth, we can prune the construction when a graph with a cycle with length less than k is constructed.

For more details on the algorithm we refer to [17].

3.2. The general case. By using our implementation of the algorithm described in Section 3.1 we generated all almost hypohamiltonian graphs with girth at least g of order n for various values of g and n . The counts of the number of almost hypo-

Algorithm 2 Construct(Graph G , int D)

```
1: if  $G$  is non-hamiltonian AND not generated before then
2:   if  $G$  contains a type A obstruction  $(W, X)$  then
3:     for every good A-edge  $e \notin E(G)$  for  $(W, X)$  for which at most one vertex has
       degree larger than  $D$  in  $G + e$  do
4:       Construct( $G + e$ ,  $D$ )
5:     end for
6:   else if  $G$  contains a vertex  $v$  of degree 2 then
7:     for every edge  $e \notin E(G)$  for which at most one vertex has degree larger than
        $D$  in  $G + e$  do
8:       Construct( $G + e$ ,  $D$ )
9:     end for
10:  else if  $G$  contains a type C obstruction  $(W, X, v)$  then
11:    for every good C-edge  $e \notin E(G)$  for  $(W, X, v)$  for which at most one vertex
      has degree larger than  $D$  in  $G + e$  do
12:      Construct( $G + e$ ,  $D$ )
13:    end for
14:  else if  $G$  contains a vertex  $v$  of degree 3 which is part of a triangle then
15:    for every edge  $e \notin E(G)$  which contains  $v$  as an endpoint for which at most
      one vertex has degree larger than  $D$  in  $G + e$  do
16:      Construct( $G + e$ ,  $D$ )
17:    end for
18:  else if  $G$  contains a type B obstruction  $(W, X)$  then
19:    for every good B-edge  $e \notin E(G)$  for  $(W, X)$  for which at most one vertex has
      degree larger than  $D$  in  $G + e$  do
20:      Construct( $G + e$ ,  $D$ )
21:    end for
22:  else
23:    if  $G$  is almost hypohamiltonian then
24:      add  $G$  to the list  $\mathcal{H}$ 
25:    end if
26:    for every edge  $e \notin E(G)$  for which at most one vertex has degree larger than
       $D$  in  $G + e$  do
27:      Construct( $G + e$ ,  $D$ )
28:    end for
29:  end if
30: end if
```

hamiltonian graphs can be found in Table 1. Therein, we also mention the number of almost hypohamiltonian graphs whose exceptional vertex is cubic—these are of special importance due to the fact that two such graphs can be combined to form a hypohamiltonian graph, see [32, Theorem 3] for details. In each case we went as far as computationally possible.

Note that we could not determine the complete set of all almost hypohamiltonian graphs with 24 vertices and girth 5—however, we succeeded in obtaining a sample, which was essential to complete the proof of Theorem 10. Combining our findings with the second author’s [32, Theorem 2], we have the following, settling [32, Problem 2].

THEOREM 10. *An almost hypohamiltonian graph of order n exists iff $n \geq 17$. The*

Order	# alm hypoham.	$g \geq 4$	$g \geq 5$	$g \geq 6$
0 – 16	0	0	0	0
17	2 (1)	2 (1)	2 (1)	0
18	2 (1)	2 (1)	1 (1)	0
19	?	27 (11)	4 (4)	0
20	?	?	14 (6)	0
21	?	?	27 (18)	0
22	?	?	133 (86)	0
23	?	?	404 (161)	0
24	?	?	≥ 68	0
25	?	?	?	0
26	?	?	?	0

Table 1: The number of almost hypohamiltonian graphs. The columns with a header of the form $g \geq k$ contain the number of almost hypohamiltonian graphs with girth at least k . In parentheses the number of almost hypohamiltonian graphs which have a cubic exceptional vertex are given.

smallest almost hypohamiltonian graph of girth 4 (girth 5) has order 18 (order 17).

Hypohamiltonian graphs of order n exist iff $n \in \{10, 13, 15, 16\}$ or $n \geq 18$, see [2]. The smallest hypohamiltonian graph of girth 4 (girth 5) has order 18 (order 10); for details we refer to [17]. No almost hypohamiltonian graphs of girths other than 4 or 5 are known, but it seems reasonable to believe that adapting a technique of Thomassen [27] will provide examples of girth 3. We know of the existence of hypohamiltonian graphs of girth g for every $g \in \{3, \dots, 7\}$, see [17], while no hypohamiltonian graphs of girth greater than 7 are known. (The smallest hypohamiltonian graph of girth 7 is Coxeter’s graph.) In Figure 2 the two smallest almost hypohamiltonian graphs (each of order 17) are depicted—both have girth 5. Figure 3 shows the smallest almost hypohamiltonian graph of girth 4 (which has order 18).

All graphs from Table 1 can also be downloaded from the *House of Graphs* [4] at <http://hog.grinvin.org/AlmostHypohamiltonian> and also be inspected in the database of interesting graphs by searching for the keywords “almost hypohamiltonian”.




Fig. 2: The almost hypohamiltonian graph of order 17 from [32]. Its exceptional vertex is w . By deleting the edge vw , we obtain the smallest almost hypohamiltonian graph both in terms of order and size.

Our generator for almost hypohamiltonian graphs was obtained by extending the code of our generator for hypohamiltonian graphs from [17]. In that paper we de-




Fig. 3: The smallest almost hypohamiltonian graph of girth 4. It has 18 vertices. Its exceptional vertex is w .

scribe how we extensively tested the correctness of our implementation. We also used multiple independent programs to test hamiltonicity and almost hypohamiltonicity—one of those programs was kindly provided to us by Gunnar Brinkmann—and in each case the results were in complete agreement. Furthermore, the source code of our new generator can be downloaded and inspected at [16].

3.3. The cubic case. The algorithm described in Section 3.1 can also be used to only generate cubic almost hypohamiltonian graphs, but it is much more efficient to use a generator for cubic graphs and test which graphs are almost hypohamiltonian as a filter.

We used the program *snarkhunter* [7, 5] to generate all cubic graphs with girth at least 4 up to 32 vertices and tested them for almost hypohamiltonicity. (Note that by Lemma 8 cubic almost hypohamiltonian graphs must have girth at least 4.) The results are presented in Table 2. The smallest cubic almost hypohamiltonian graphs of girth 4 and 5 are shown in Figure 4.

We also used the complete lists of snarks from [6] to determine all almost hypohamiltonian snarks up to 36 vertices. A *snark* is a cyclically 4-edge-connected cubic graph with girth at least 5 which is not 3-edge-colourable. The number of almost hypohamiltonian snarks is listed in the last column of Table 2. The sudden increase at 34 vertices is striking and also occurs for hypohamiltonian snarks as described in [6].

All almost hypohamiltonian graphs from Table 2 can be downloaded from the *House of Graphs* [4] at <http://hog.grinvin.org/AlmostHypohamiltonian> and also be inspected in the database of interesting graphs by searching for the keywords “almost hypohamiltonian”.

Using *snarkhunter* we also verified that there are no cubic almost hypohamiltonian graphs with girth 6 up to at least 36 vertices.

By combining the results from Table 2 with the results on non-hamiltonian cubic graphs from [17] we obtain the following bounds for cubic almost hypohamiltonian graphs.

THEOREM 11. *Let c_g denote the order of the smallest cubic almost hypohamiltonian graph with girth g . We have*

$$c_4 = 28, \quad c_5 = 26, \quad c_6 \geq 38, \quad c_7 \geq 44, \quad c_8 \geq 50, \quad \text{and} \quad c_9 \geq 66.$$




Fig. 4: Figure 4a depicts one of the ten smallest cubic almost hypohamiltonian graphs of girth 5. It has 26 vertices. Figure 4b shows one of the four smallest cubic almost hypohamiltonian graphs of girth 4. It has 28 vertices. In both figures the exceptional vertex is labelled w .

Order	$g \geq 4$	Non-ham. and $g \geq 4$	Alm. hypo.	Alm. hypo. and $g \geq 5$	Alm. hypo. and $g \geq 6$	Alm. hypo. snark
6	1	0	0	0	0	0
8	2	0	0	0	0	0
10	6	1	0	0	0	0
12	22	0	0	0	0	0
14	110	2	0	0	0	0
16	792	8	0	0	0	0
18	7 805	59	0	0	0	0
20	97 546	425	0	0	0	0
22	1 435 720	3 862	0	0	0	0
24	23 780 814	41 293	0	0	0	0
26	432 757 568	518 159	10	10	0	10
28	8 542 471 494	7 398 734	6	2	0	2
30	181 492 137 812	117 963 348	25	12	0	11
32	4 127 077 143 862	2 069 516 990	74	4	0	0
34	?	?	?	?	?	6 253
36	?	?	?	?	?	2 243

Table 2: Counts of almost hypohamiltonian graphs among cubic graphs. g stands for girth. Note that due to Lemma 8, the girth of a cubic almost hypohamiltonian graph is at least 4. The last column denotes the number of almost hypohamiltonian snarks.

In comparison, if we denote by c'_g the smallest cubic hypohamiltonian graph with girth g , we have the following. (See [17] for more details.)

$$c'_4 = 24, \quad c'_5 = 10, \quad c'_6 = c'_7 = 28, \quad c'_8 \geq 50, \quad \text{and} \quad c'_9 \geq 66.$$

3.4. The planar case. Until recently, the smallest known planar almost hypohamiltonian graph had order 39, see [32]. [32, Problem 3] and [33, Problem 6] ask for the smallest order of a planar almost hypohamiltonian graph, and the smallest number n_0 such that there exists a planar almost hypohamiltonian graph of order n for every $n \geq n_0$. Although we are not able to fully settle the questions, we can report progress on both the lower and upper bounds of the order of the smallest planar

almost hypohamiltonian graph, as well as an improvement of n_0 .

We shall strengthen the second author's [32, Theorem 1] and [32, Theorem 4]. Although Wiener recently has shown [31] the existence of a planar almost hypohamiltonian graph of order 31 (which constitutes the currently best upper bound for the order of the smallest planar almost hypohamiltonian graph), the planar almost hypohamiltonian graphs we will construct in Theorem 18—the smallest among them has 36 vertices—have a cubic exceptional vertex, while the exceptional vertex in Wiener's 31-vertex graph has degree 4. Cubic exceptional vertices are of particular interest due to the fact that we can combine two almost hypohamiltonian graphs, each having a cubic exceptional vertex, into a hypohamiltonian graph, see [32, Theorem 3]—no method is known to do this if the exceptional vertex is non-cubic.

Let us first focus on the lower bound for the order of the smallest planar almost hypohamiltonian graph. Since the algorithm for generating all almost hypohamiltonian graphs presented in Section 3.1 only adds edges and never removes any vertices or edges, all graphs obtained by the algorithm from a non-planar graph will remain non-planar. So in case we only want to generate planar almost hypohamiltonian graphs, we can prune the construction when a non-planar graph is constructed. We used Boyer and Myrvold's algorithm [3] to test if a graph is planar.

Following this approach, we showed that the smallest planar almost hypohamiltonian graph has at least 22 vertices and that the smallest planar almost hypohamiltonian graph of girth 4 must have at least 26 vertices. For girth 5 it turned out to be more efficient to use the program *plantri* [8] instead. Using *plantri* we generated all planar 3-connected graphs with girth 5 up to 48 vertices and tested them for almost hypohamiltonicity. This yielded the following results.

THEOREM 12. *The smallest planar almost hypohamiltonian graph with girth 5 has 44 vertices. There is exactly one such graph of that order and it is shown in Figure 5. There are exactly 42 planar almost hypohamiltonian graphs with girth 5 on 47 vertices. These are the only planar almost hypohamiltonian graphs with girth 5 up to at least 48 vertices.*

The graphs from Theorem 12 can be downloaded from the database of interesting graphs at the *House of Graphs* [4] by searching for the keywords “planar almost hypohamiltonian”.

In summary, we have the following lower bounds.

COROLLARY 13. *Let $\bar{h}$ ($\bar{h}_g$) denote the order of the smallest planar almost hypohamiltonian graph (of girth g). We have*

$$\bar{h} \geq 22, \quad \bar{h}_4 \geq 26, \quad \text{and} \quad \bar{h}_5 = 44.$$

Wiener [31] showed that there exists a planar almost hypohamiltonian graph of order 31 and girth 4, so $\bar{h} \leq \bar{h}_4 \leq 31$. (No planar almost hypohamiltonian graph of girth 3 is known. Adapting a technique of Thomassen [27] could provide examples of girth 3, as mentioned earlier.) In comparison, if we denote by $\bar{h}'$ ($\bar{h}'_g$) the order of the smallest planar hypohamiltonian graph (of girth g), we have the following—more details can be found in [17].

$$23 \leq \bar{h}' \leq 40, \quad 23 \leq \bar{h}'_3 \leq 240, \quad 25 \leq \bar{h}'_4 \leq 40, \quad \text{and} \quad \bar{h}'_5 = 45.$$

We now prepare for showing that there exists a planar almost hypohamiltonian graph of order 36 whose exceptional vertex is cubic, as well as improving n_0 (defined in the first paragraph of this subsection) from 76 (see [32]) to 73. Let G be a graph




Fig. 5: The smallest planar almost hypohamiltonian graph of girth 5. It has 44 vertices. Its exceptional vertex is w .

containing a 4-cycle $v_1v_2v_3v_4 = C$. We denote by $\text{Th}(G_C)$ the graph obtained from G by deleting the edges v_1v_2 and v_3v_4 and adding a 4-cycle $v'_1v'_2v'_3v'_4$ disjoint from G , and the edges $v_iv'_i$, $1 \leq i \leq 4$, to G . (Note that the operation Th preserves planarity. It was introduced by Thomassen [30] to show that there exist infinitely many planar cubic hypohamiltonian graphs.) The statement of Lemma 1 is a slight modification of a claim of Thomassen, see [30]; we skip its proof.

LEMMA 14 (Thomassen [30]). *Let G be a planar non-hamiltonian graph containing a quadrilateral face bounded by the cycle C . Then $\text{Th}(G_C)$ is planar and non-hamiltonian.*

In a planar graph, we call a face *cubic* if all of its vertices are cubic. In [30], Thomassen also showed that given a planar hypohamiltonian graph G which contains a cubic quadrilateral face bounded by the cycle C , we have that $\text{Th}(G_C)$ is a planar hypohamiltonian graph, as well. We require a modified version of Thomassen's result.

LEMMA 15 (Zamfirescu [32]). *Let G be a planar almost hypohamiltonian graph with exceptional vertex w , and let G contain a cubic quadrilateral face bounded by the cycle C such that $w \notin V(C)$. Then $\text{Th}(G_C)$ is a planar almost hypohamiltonian graph, as well.*

Consider graphs G and H , and the cubic vertices $x \in V(G)$ and $y \in V(H)$. Denote by G_xH_y one of the graphs obtained from $G - x$ and $H - y$ by identifying the vertices in $N(x)$ with those in $N(y)$ using a bijection (so G_xH_y has $|V(G)| + |V(H)| - 5$ vertices). Thomassen [26] showed that if G and H are hypohamiltonian, then G_xH_y is hypohamiltonian, as well. Note that if G is hypohamiltonian, then G contains no triangle with a cubic vertex (as shown by Collier and Schmeichel [12, p. 196]). We will need the following lemma.

LEMMA 16 (Zamfirescu [32]). *Let G be an almost hypohamiltonian graph which contains a cubic vertex x different from the exceptional vertex w of G , and H a hypohamiltonian graph which contains a cubic vertex y . Then G_xH_y is an almost*

hypohamiltonian graph with exceptional vertex w . If G and H are planar, then so is $G_x H_y$.

A further ingredient is the following useful result of Grinberg.

THEOREM 17 (Grinberg's Criterion [18]). *Given a plane graph with a hamiltonian cycle $\mathfrak{h}$ and exactly f_i (f'_i) i -gons inside (outside) of $\mathfrak{h}$, we have*

$$\sum_{i \geq 3} (i-2)(f_i - f'_i) = 0. \quad (\dagger)$$

THEOREM 18. *There exists a planar almost hypohamiltonian graph of order n and with cubic exceptional vertex for (i) $n = 36 + 4k$, $k \geq 0$, and (ii) every $n \geq 73$.*

Proof. (i) We first show that the graph G from Figure 6a is almost hypohamiltonian. By Grinberg's Criterion, G is non-hamiltonian, since all faces of G are pentagons with exactly one exception, which is a quadrilateral—now the left-hand side of $(\dagger)$ cannot vanish. $G - w$ is non-hamiltonian, as well: denote by $Q = a'b'c'd'$ the quadrilateral in G , by a' the cubic vertex of Q , and be w the exceptional vertex of G , as shown in Figure 6a. Assume that there exists a hamiltonian cycle $\mathfrak{h}$ in $G - w$. By Grinberg's Criterion [18], in $G - w$, Q and the nonagon N must lie on the same side of $\mathfrak{h}$. But this is impossible, as $a' \in V(\mathfrak{h})$ and thus, Q and N lie on different sides of $\mathfrak{h}$. Hence, $G - w$ is non-hamiltonian.




Fig. 6: G and G' (i.e. Figure 6a and 6b, respectively), two planar almost hypohamiltonian graphs used in the proof of Theorem 18. Their respective cubic exceptional vertices are labelled w .

We skip the straightforward verification that for every vertex $v \neq w$, the graph $G - v$ is hamiltonian. Thus, we have shown that there exists a planar almost hypohamiltonian graph of order 36 whose exceptional vertex is cubic. Note that we cannot apply Lemma 15 to G since $a'b'c'd'$ is not a cubic quadrilateral, thus we require the following claim.

Claim. *The graph G' from Figure 6b is planar and almost hypohamiltonian with exceptional vertex w .*

Proof of the Claim. Since $G' = \text{Th}(G_Q)$, by Lemma 14 G' is planar and non-hamiltonian. Let $\mathfrak{h}$ be a hamiltonian cycle in $G' - w$. Certainly, $aa'd' \subset \mathfrak{h}$. If

$dd' \in E(\mathfrak{h})$, then $b'baa'd'dcc' \subset \mathfrak{h}$, which replaced with $b'a'd'd'c'$ gives a hamiltonian cycle in $G - w$, a contradiction, so $dd' \notin E(\mathfrak{h})$. Hence $a'adcbb' \subset \mathfrak{h}$, which replaced with $a'b'$ gives a hamiltonian cycle in $G - w$, once more a contradiction.

It is quickly verified that there exists a hamiltonian cycle $\mathfrak{h}$ in $G - a'$ containing the path $b'c'd'$. Deleting from $\mathfrak{h}$ the edge $c'd'$ we obtain the path $\mathfrak{p}$ in G' . Now $\mathfrak{p} \cup d'dabcc'$ is a hamiltonian cycle in $G' - a'$. Finding hamiltonian cycles in $G' - b'$, $G' - c'$, and $G' - d'$ works in a similar (and in fact even simpler) fashion. $\mathfrak{p} \cup d'a'adcc'b'$ is a hamiltonian cycle in $G' - b$, and $\mathfrak{p} \cup d'a'abcc'b'$ is a hamiltonian cycle in $G' - d$. It is easy to see that there exists a hamiltonian cycle $\mathfrak{h}'$ in $G - d'$ containing the edge $b'a'$. Deleting this edge from $\mathfrak{h}'$ we obtain the path $\mathfrak{p}$ in G' . Now $\mathfrak{p} \cup b'bcdd'a'$ is a hamiltonian cycle in $G' - a$, and $\mathfrak{p} \cup b'badd'a'$ is a hamiltonian cycle in $G' - c$.

It remains to show that $G' - v'$ is hamiltonian for every $v' \in V(G') \setminus \{w, a, a', b, b', c, c', d, d'\}$. Let v be the vertex in G corresponding to v' in G' . Let $\mathfrak{h}''$ be a hamiltonian cycle in $G - v$. If $a'b' \in E(\mathfrak{h}'')$, replace $a'b'$ with $a'adcbb'$. If $a'b' \notin E(\mathfrak{h}'')$, then necessarily $a'd' \in E(\mathfrak{h}'')$, which we replace with $a'abcdd'$. In both cases we have obtained a hamiltonian cycle in $G' - v'$, which finishes the proof of the Claim.

Using the Claim and iterating Lemma 15 ad infinitum, the proof of (i) is complete.

(ii) In [22], it was shown that for every $n \geq 42$ there exists a planar hypohamiltonian graph of order n , which we will call H^n . For arbitrary but fixed $n \geq 42$, let $x \in V(H^n)$ be cubic (Thomassen showed that every planar hypohamiltonian graph contains a cubic vertex [29]), and let $y \in V(G)$ be cubic, as well. (Where G is the graph shown in Figure 6a.) Applying Lemma 16, we obtain that $H_x^n G_y$ is an almost hypohamiltonian graph with a cubic exceptional vertex. $\square$

By applying [32, Theorem 3], Theorem 18 (i) yields an alternative proof of the fact that infinitely many planar hypohamiltonian graphs exist. (This was first shown by Thomassen [28]. Chvátal [11] had asked whether the statement is true, and Grünbaum [19, p. 37] had conjectured that it is not.)

3.5. The planar cubic case. Using the program *plantri* [8] we generated all planar cyclically 4-edge-connected cubic graphs with girth at least 4 up to 52 vertices and tested them for almost hypohamiltonicity. No such graphs were found, so we have:

THEOREM 19. *The smallest planar cubic almost hypohamiltonian graph has at least 54 vertices.*

With *plantri* we also generated all planar cyclically 4-edge-connected cubic graphs with girth 5 up to 78 vertices. This yielded the following results. These results were independently obtained by McKay (private communication) but were not published.

THEOREM 20. *The smallest planar cubic almost hypohamiltonian graph of girth 5 has 68 vertices. There are exactly three such graphs of that order and they are shown in Figure 7. There are exactly 81 planar cubic almost hypohamiltonian graphs with girth 5 on 74 vertices. These are the only planar cubic almost hypohamiltonian graphs with girth 5 up to at least 78 vertices.*

The graphs from Theorem 20 can be downloaded from the database of interesting graphs at the *House of Graphs* [4] by searching for the keywords “planar cubic almost hypohamiltonian”.

The smallest planar cubic hypohamiltonian graph of girth 5 has order 76, as proven by McKay [23]. In [17], we showed that there exists exactly one such graph of order 78. Interestingly, McKay’s three extremal graphs of order 76 have trivial auto-




Fig. 7: All three smallest planar cubic almost hypohamiltonian graphs of girth 5. Each has 68 vertices. In each figure the exceptional vertex is labelled w .

morphism group, whereas our 78-vertex example has D_{3h} symmetry (as an abstract group, this is the dihedral group of order 12).

Acknowledgements. Most computations for this work were carried out using the Stevin Supercomputer Infrastructure at Ghent University. We also would like to thank Gunnar Brinkmann for providing us with an independent program for testing almost hypohamiltonicity and Brendan McKay for his advice on *plantri*.

REFERENCES

- [1] G. M. ADEL'SON-VEL'SKIĬ AND V. K. TITOV, *On edge 4-chromatic cubic graphs (in Russian)*, in Proc. Seminar of 1971 at Moscow Univ., Voprosy Kibernetiki [vol. not numbered], 1973, pp. 5–14.
- [2] R. E. L. ALDRED, B. D. MCKAY, AND N. C. WORMALD, *Small hypohamiltonian graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing, 23 (1997), pp. 143–152.
- [3] J. M. BOYER AND W. J. MYRVOLD, *On the Cutting Edge: Simplified $O(n)$ Planarity by Edge Addition*, Journal of Graph Algorithms and Applications, 8 (2004), pp. 241–273.
- [4] G. BRINKMANN, K. COOLSAET, J. GOEDGEBEUR, AND H. MÉLOT, *House of Graphs: a database of interesting graphs*, Discrete Applied Mathematics, 161 (2013), pp. 311–314. Available at <http://hog.grinvin.org/>.
- [5] G. BRINKMANN AND J. GOEDGEBEUR, *Generation of cubic graphs and snarks with large girth*, arXiv preprint arXiv:1512.05944, (2015).
- [6] G. BRINKMANN, J. GOEDGEBEUR, J. HÄGGLUND, AND K. MARKSTRÖM, *Generation and properties of snarks*, Journal of Combinatorial Theory, Series B, 103 (2013), pp. 468–488.
- [7] G. BRINKMANN, J. GOEDGEBEUR, AND B. D. MCKAY, *Generation of cubic graphs*, Discrete Mathematics and Theoretical Computer Science, 13 (2011), pp. 69–80.
- [8] G. BRINKMANN AND B. D. MCKAY, *Fast generation of planar graphs*, MATCH Commun. Math. Comput. Chem., 58 (2007), pp. 323–357.
- [9] A. CAVICCHIOLI, T. E. MURGOLO, B. RUINI, AND F. SPAGGIARI, *Special classes of snarks*, Acta Applicandae Mathematica, 76 (2003), pp. 57–88.
- [10] G. CHARTRAND, S. F. KAPOOR, AND D. R. LICK, *n -hamiltonian graphs*, Journal of Combinatorial Theory, 9 (1970), pp. 308–312.
- [11] V. CHVÁTAL, *Flip-flops in hypohamiltonian graphs*, Canadian Mathematical Bulletin, 16 (1973), pp. 33–41.
- [12] J. B. COLLIER AND E. F. SCHMEICHEL, *Systematic searches for hypohamiltonian graphs*, Networks, 8 (1978), pp. 193–200.
- [13] R. DIESTEL, *Graph Theory*, Graduate Texts in Mathematics, 173 (2010).
- [14] S. FIORINI, *Hypohamiltonian snarks*, in Graphs and Other Combinatorial Topics (ed.: M. Fiedler), vol. 59, Teubner, 1983, pp. 70–75.
- [15] J. GOEDGEBEUR, E. STEFFEN, AND C. T. ZAMFIRESCU, *On hypohamiltonian snarks and*

- fiorini's theorem*, In preparation.
- [16] J. GOEDGEBEUR AND C. T. ZAMFIRESCU, *Homepage of genhypohamiltonian*: <http://caagt.ugent.be/hypoham/>.
 - [17] J. GOEDGEBEUR AND C. T. ZAMFIRESCU, *Improved bounds for hypohamiltonian graphs*, arXiv preprint arXiv:1602.07171, (2016).
 - [18] E. J. GRINBERG, *Plane homogeneous graphs of degree three without hamiltonian circuits (in Russian)*, Latvian Math. Yearbook, 4 (1968), pp. 51–58.
 - [19] B. GRÜNBAUM, *Vertices missed by longest paths or circuits*, Journal of Combinatorial Theory, Series A, 17 (1974), pp. 31–38.
 - [20] D. A. HOLTON AND J. SHEEHAN, *The Petersen graph, chapter 7: Hypohamiltonian graphs*, 1993.
 - [21] R. ISAACS, *Infinite families of nontrivial trivalent graphs which are not Tait colorable*, American Mathematical Monthly, 82 (1975), pp. 221–239.
 - [22] M. JOOYANDEH, B. D. MCKAY, P. R. J. ÖSTERGÅRD, V. H. PETTERSSON, AND C. T. ZAMFIRESCU, *Planar hypohamiltonian graphs on 40 vertices*, To appear in: Journal of Graph Theory, (2016).
 - [23] B. D. MCKAY, *Hypohamiltonian planar cubic graphs with girth 5*, To appear in: Journal of Graph Theory, (2016).
 - [24] Z. SKUPIEŃ, *Multi-compositions in exponential counting of hypohamiltonian snarks*, in Convexity and Discrete Geometry Including Graph Theory, Springer Proc. in Math. & Stat. (eds.: K. Adiprasito, I. Bárány, C. Vilcu), vol. 148, Springer, 2016, pp. 47–58.
 - [25] R. SOUSSELIER, *Problème no. 29: Le cercle des irascibles (in French)*, Revue Française de Recherches Opérationnelle, 7 (1963), pp. 405–406.
 - [26] C. THOMASSEN, *Hypohamiltonian and hypotraceable graphs*, Discrete Mathematics, 9 (1974), pp. 91–96.
 - [27] C. THOMASSEN, *On hypohamiltonian graphs*, Discrete Mathematics, 10 (1974), pp. 383–390.
 - [28] C. THOMASSEN, *Planar and infinite hypohamiltonian and hypotraceable graphs*, Discrete Mathematics, 14 (1976), pp. 377–389.
 - [29] C. THOMASSEN, *Hypohamiltonian graphs and digraphs*, in Theory and Applications of Graphs, Springer, 1978, pp. 557–571.
 - [30] C. THOMASSEN, *Planar cubic hypohamiltonian and hypotraceable graphs*, Journal of Combinatorial Theory, Series B, 30 (1981), pp. 36–44.
 - [31] G. WIENER, *New constructions of hypohamiltonian and hypotraceable graphs*, In preparation.
 - [32] C. T. ZAMFIRESCU, *On hypohamiltonian and almost hypohamiltonian graphs*, Journal of Graph Theory, 79 (2015), pp. 63–81.
 - [33] C. T. ZAMFIRESCU, *Seven problems on hypohamiltonian and almost hypohamiltonian graphs*, in Convexity and Discrete Geometry Including Graph Theory, Springer Proc. in Math. & Stat. (eds.: K. Adiprasito, I. Bárány, C. Vilcu), vol. 148, Springer, 2016, pp. 253–255.